

Prospective Risk-Adjusted [¹⁸F]Fluorodeoxyglucose Positron Emission Tomography and Computed Tomography Assessment of Radiation Response in Head and Neck Cancer

Benjamin J. Moeller, Vishal Rana, Blake A. Cannon, Michelle D. Williams, Erich M. Sturgis, Lawrence E. Ginsberg, Homer A. Macapinlac, J. Jack Lee, K. Kian Ang, K.S. Clifford Chao, Gregory M. Chronowski, Steven J. Frank, William H. Morrison, David I. Rosenthal, Randal S. Weber, Adam S. Garden, Scott M. Lippman, and David L. Schwartz

From the Departments of Radiation Oncology, Pathology, Head and Neck Surgery, Diagnostic Imaging, Nuclear Medicine, Biostatistics, and Thoracic/Head and Neck Medical Oncology, The University of Texas M. D. Anderson Cancer Center, Houston, TX.

Submitted August 15, 2008; accepted December 12, 2008; published online ahead of print at www.jco.org on March 30, 2009.

Supported by The University of Texas M.D. Anderson Cancer Center Specialized Programs of Research Excellence (SPORE) in Head and Neck Grant No. CA97007.

Authors' disclosures of potential conflicts of interest and author contributions are found at the end of this article.

Corresponding author: David L. Schwartz, MD, Department of Radiation Oncology, The University of Texas M. D. Anderson Cancer Center, Box 97, 1515 Holcombe Blvd, Houston, TX, 77030; e-mail: docdls@mdanderson.org.

The Appendix is included in the full-text version of this article, available online at www.jco.org. It is not included in the PDF version (via Adobe® Reader®).

© 2009 by American Society of Clinical Oncology

0732-183X/09/2715-2509/\$20.00

DOI: 10.1200/JCO.2008.19.3300

A B S T R A C T

Purpose

[¹⁸F]Fluorodeoxyglucose positron emission tomography (FDG-PET)/computed tomography (CT) imaging may improve assessment of radiation response in patients with head and neck cancer, but it is not yet known for which patients this is most useful. We conducted a prospective trial to identify patient populations likely to benefit from the addition of functional imaging to the assessment of radiotherapy response.

Patients and Methods

Ninety-eight patients with locally advanced cancer of the oropharynx, larynx, or hypopharynx were prospectively enrolled and treated with primary radiotherapy, with or without chemotherapy. Patients underwent FDG-PET/CT and contrast-enhanced CT imaging 8 weeks after completion of treatment. Functional and anatomic imaging response was correlated with clinical and pathologic response. Imaging accuracy was then compared between imaging modalities.

Results

Although postradiation maximum standard uptake values were significantly higher in nonresponders compared with responders, the positive and negative predictive values of FDG-PET/CT scanning were similar to those for CT alone in the unselected study population. Subset analyses revealed that FDG-PET/CT outperformed CT alone in response assessment for patients at high risk for treatment failure (those with human papillomavirus [HPV]–negative disease, nonoropharyngeal primaries, or history of tobacco use). No benefit to FDG-PET/CT was seen for low-risk patients lacking these features.

Conclusion

These data do not support the broad application of FDG-PET/CT for radiation response assessment in unselected head and neck cancer patients. However, FDG-PET/CT may be the imaging modality of choice for patients with highest risk disease, particularly those with HPV-negative tumors. Optimal timing of FDG-PET/CT imaging after radiotherapy merits further investigation.

J Clin Oncol 27:2509-2515. © 2009 by American Society of Clinical Oncology

INTRODUCTION

Primary radiotherapy, with or without concurrent chemotherapy, for organ-sparing treatment of head and neck squamous cell carcinoma (HNSCC) has advanced markedly over the last decade. However, there remains room for improvement, with local control rates as low as 50% in recent phase III trials.¹⁻³ For patients with persistent disease after radiation, consolidative neck dissection and/or surgical salvage of primary disease potentially improve survival.⁴ However, such

surgery carries significant risk for morbidity and mortality.⁵⁻⁷ This underscores the importance of correctly identifying those patients in need of salvage therapy.⁸

The ideal method of treatment response assessment would noninvasively achieve accurate prediction of residual viable tumor after radiotherapy. Clinical examination alone is inadequate for response assessment, providing accuracy less than 50%.^{9,10} Postradiation computed tomography (CT) imaging yields higher sensitivity but has been criticized for lacking specificity.^{11,12}

Recent studies have proposed that functional imaging, specifically [^{18}F]fluorodeoxyglucose positron emission tomography (FDG-PET)/CT, may improve the accuracy of radiation response assessment for HNSCC.^{8,13-22} These studies are limited by their retrospective design, and consequently, the role of FDG-PET/CT in radiation response assessment remains undefined.

We sought to more definitively address this issue by conducting a prospective trial designed to compare the accuracy of radiation response assessment by FDG-PET/CT and contrast-enhanced CT and define patient subsets likely to derive maximal benefit from the addition of FDG-PET/CT imaging to conventional response assessment.

PATIENTS AND METHODS

Patients

Between November 2005 and May 2007, 107 consecutive patients with locally advanced HNSCC scheduled to receive definitive radiation with or without concurrent systemic therapy were screened for enrollment onto this prospective institutional review board–approved trial. Adult patients were eligible for inclusion if they had biopsy-proven American Joint Commission on Cancer (AJCC) version 6 stage III to IVB squamous cell carcinoma of the oropharynx, hypopharynx, or larynx. Nine screened patients were ineligible for inclusion; two withdrew consent, five had disease removed surgically before radiotherapy, and two had FDG-PET/CT scans performed earlier than as stipulated by protocol, leaving an enrolled study cohort of 98 patients.

Demographic and clinical covariates were directly tabulated from each patient's clinical chart. A threshold level of 10 pack-years of use was chosen to distinguish tobacco users from nonusers because this level of exposure is correlated with increased risk of developing HNSCC.²³

Radiotherapy and Systemic Treatment

Patients were treated with intensity-modulated radiotherapy as described previously.^{24,25} Prescription doses ranged from 66 to 70 Gy in 32 to 35 fractions. Intensity-modulated radiotherapy was delivered via a step-and-shoot, multileaf collimation through a static treatment gantry. Treatment planning was performed with a Pinnacle³ system (version 6.2b or later; Philips Medical Systems, Andover, MA). Seventy-eight patients received concurrent systemic therapy with radiation.

FDG-PET/CT Imaging

Two FDG-PET/CT scans were scheduled for each patient; the first was scheduled within 4 weeks before starting therapy, and the second was scheduled as near as possible to the eighth week after completion of treatment. Deviations from the 8-week target were allowed, as long as imaging was performed between 5 and 12 weeks after radiotherapy. Details of our functional imaging protocol are provided in the Appendix (online only). The intensity of metabolic activity within foci of increased FDG uptake in disease sites was analyzed on the FDG-PET images using a semiautomated vendor-provided tool (GE Advantage Workstation; GE Medical Systems, Milwaukee, WI). A single clinician (V.R.) calculated maximum standardized uptake values (SUV_{max}) from a volume of interest completely encompassing each site of FDG accumulation. For patients with multiple sites of lymphadenopathy, nodal SUV_{max} was calculated on each FDG-PET/CT scan for the node with the highest initial SUV_{max} . FDG-PET/CT scans were interpreted with knowledge of contrast-enhanced CT scans.

CT Imaging

Two contrast-enhanced head and neck CT scans were scheduled for each patient; the first was scheduled before starting therapy, and the second was scheduled as near as possible to the eighth week after treatment. All images were interpreted by a board-certified neuroradiologist (L.E.G.). Interpretation

of conventional CT imaging was blinded to FDG-PET/CT results. For further details of the anatomic imaging protocol, refer to the Appendix.

Post-Treatment Surveillance and Surgical Management

Patients returned for routine post-treatment surveillance 8 weeks after treatment for baseline assessment and then every 3 to 4 months for the first 2 years for clinical examination and serial contrast-enhanced CT imaging of the head and neck. Chest x-rays were performed annually.

A standardized clinical pathway was followed for postradiotherapy neck management (see Appendix). Consolidative neck dissections were performed at the discretion of the treating physicians and as indicated by findings from all post-treatment studies including, but not limited to, FDG-PET/CT. Patients with no radiographic or clinical evidence for residual adenopathy were observed expectantly. Patients found by clinical or radiographic evaluation to have residual or recurrent primary disease underwent salvage resection.

Histology was the gold standard for defining nonresponse to therapy for patients who underwent surgical consolidation. For patients observed clinically, the gold standard was any locoregional failure detected by physical examination (including fiberoptic pharyngolaryngoscopy) or radiographic imaging (CT, PET/CT, and/or ultrasound) within 6 months of completing radiotherapy.

Pathologic Tissue Assessment

Twenty-two patients underwent consolidative neck dissection or salvage resection of primary disease at a median of 85 days after completion of radiotherapy. All neck dissection tissue specimens were evaluated and step sectioned by the same board-certified pathologist (M.D.W.). For further details of the step-sectioning protocol, refer to the Appendix.

Human Papillomavirus Detection

When tissue was available, polymerase chain reaction (PCR), in situ hybridization (ISH), and immunohistochemistry were run on paraffin-embedded tumor samples to detect the presence or absence of human papillomavirus (HPV). See the Appendix for further details. Samples were regarded as positive if HPV was detected by either PCR or ISH. Discrepancies between methods were rare and were resolved by p16 immunohistochemistry; positive staining for p16, with HPV DNA detected by either PCR or ISH, was coded as positive for HPV.

Data Analysis

Locoregional disease control, distant metastasis-free survival, disease-specific survival, and overall survival from the date of treatment completion were estimated using the Kaplan-Meier method. Patients were separated into two groups, complete responders and nonresponders, based on clinical and pathologic response assessment as described earlier. Multivariate Cox regression analysis (WinStat; <http://www.winstat.com/>) was used to analyze covariates for correlation with failure (significance defined as $P < .05$). SUV_{max} values were compared between the groups using a two-tailed t test (significance defined as $P < .05$). Receiver operating characteristic (ROC) curve analysis was performed for the postradiation FDG-PET/CT data sets, separately for primary and nodal disease. Polynomial curves were fit to the ROC data and solved for a slope of 1 to identify SUV_{max} thresholds with maximum accuracy for predicting failure. These values were then applied to the raw postradiation FDG-PET/CT data to calculate the accuracy of predicting response (see Appendix).

RESULTS

Patient Characteristics

The median follow-up time for all patients enrolled onto this study was 92 weeks (range, 37 to 122 weeks). Two years after the completion of radiotherapy, the predicted overall survival rate for the enrolled cohort was 89.1%, disease-specific survival rate was 90.3%, locoregional control rate was 88.2%, and distant




Fig 1. Maximum standardized uptake value (SUV_{max}) distribution for responders and nonresponders. Histogram plots of (A and B) preradiation and (C and D) postradiation SUV_{max} values are shown for (A and C) primary tumors and (B and D) nodes. An adjusted scale is shown for postradiation (E) primary tumors and (F) nodes for ease of viewing.

metastasis-free survival rate was 93.1%. The baseline demographic and clinical characteristics of the enrolled patients were typical for locally advanced HNSCC patients treated at our institution (Appendix Table A1, online only). The majority of patients were nonsmoking white males with stage IV oropharyngeal squamous cell carcinoma treated with 70 Gy over 6 weeks with concurrent weekly cisplatin.

Six patients were censored after experiencing distant failure and initiating systemic therapy before developing locoregional recurrence. This left 92 patients available for further analysis of locoregional disease response. These patients underwent surgical or clinical evaluation of treatment response at the discretion of the treating physician. Eighty patients responded to radiotherapy, and 12 were nonresponders (Appendix Fig A1, online only). Multivariate Cox regression analysis was performed to examine whether baseline clinical or demographic data predicted for response to therapy.

Advanced T stage and a history of significant tobacco use were both significant predictors of failure ($P = .045$ and $P = .038$, respectively).

Correlating SUV_{max} With Treatment Response

We first examined whether SUV_{max} values correlated with response to treatment (Fig 1). There were no significant differences between responders and nonresponders in baseline SUV_{max} values for primary tumors or nodes (Figs 1A and 1B). However, after radiation, SUV_{max} values were significantly higher in nonresponders compared with responders for both primary tumors and nodes ($P < .001$; Figs 1C to 1F). We also examined whether relative or absolute change in SUV_{max} from the pre- to postradiation FDG-PET/CT scans correlated with response (Appendix Table A2, online only). The relative change in SUV_{max} for primary tumors was the only other parameter significantly associated with response to treatment ($P < .001$).

Accuracy of Post-Treatment SUV_{max} for Predicting Treatment Response

Because absolute postradiation SUV_{max} was the sole factor correlating with response for both primary tumors and nodal disease, we focused the remainder of our analyses on these data. ROC curve analysis was performed on the postradiation FDG-PET/CT data set to define threshold SUV_{max} values with highest accuracy for radiation response assessment (Fig 2). We identified 6.5 and 2.8 as threshold SUV_{max} values with maximal accuracy for predicting failure in primary tumors and nodes, respectively. Clinical interpretation of each FDG-PET/CT scan agreed with the results of SUV_{max} analysis for all patients on study.

Using these threshold values, the sensitivity, specificity, positive predictive value, and negative predictive value were each calculated for the postradiation FDG-PET/CT data set. For comparison, these values were also calculated for the CT data set, using complete radiographic response as the predictor for clinical cure and all other radiographic end points (partial response, stable disease, and progressive disease) as predictors for failure. Overall, the accuracy of FDG-PET/CT was similar to that of CT (Table 1). When compared with CT alone, FDG-PET/CT had slightly higher specificity and positive predictive value at a cost of lower sensitivity and negative predictive value.

Risk Stratification and FDG-PET/CT Accuracy

The positive predictive value of FDG-PET/CT in our patient cohort was relatively low; it performed better than chance only in predicting residual primary disease and then only by a small margin. The positive predictive value of any test can be improved by limiting its use to patients with a high incidence of events. Therefore, we reasoned that the accuracy of FDG-PET/CT in radiation response assessment could be maximized by applying it to patients at highest risk for treatment failure.

We divided our patients into low-risk and high-risk groups by the following factors correlating with a lack of response to treatment: T stage (0 to 2 ν 3 to 4), N stage (0 to 2a ν 2b to 3), overall




Fig 2. [18 F]fluorodeoxyglucose positron emission tomography (FDG-PET)/computed tomography (CT) receiver operating characteristic curves are shown (primary tumors = —; nodal disease = - - - -). For comparison, the sensitivity/specificity of CT alone is plotted for primary tumors (○) and nodal disease (□), as well.

Table 1. Accuracy of FDG-PET/CT ν CT

Factor	Primary		Node	
	PET/CT	CT	PET/CT	CT
True positive, No.	7	8	6	7
False negative, No.	3	2	2	1
True negative, No.	74	71	51	44
False positive, No.	5	8	16	23
Sensitivity, %	70.0	80.0	75.0	87.5
NPV, %	96.1	97.3	96.2	97.8
Specificity, %	93.7	89.9	76.1	65.7
PPV, %	58.3	50.0	27.3	23.3

NOTE. Raw numbers of true and false negatives and positives are shown. From these numbers were derived sensitivity, specificity, PPV, and NPV. All values are tabulated for FDG-PET/CT and CT by primary and nodal disease.

Abbreviations: FDG, [18 F]fluorodeoxyglucose; PET, positron emission tomography; CT, computed tomography; NPV, negative predictive value; PPV, positive predictive value.

American Joint Commission on Cancer stage (II to III ν IV), primary tumor site (oropharyngeal ν nonoropharyngeal), smoking history (user ν nonuser), and tumor HPV status (positive ν negative). When possible, groupings were designed to divide patients into equal-sized subgroups.

The positive predictive value of FDG-PET/CT did not surpass 50% for any patient subset separated by stage (T, N, or overall). However, this objective was reached for patients with nonoropharyngeal primaries, positive smoking histories, and HPV-negative tumors (Fig 3). The greatest advantage in positive predictive value was seen by dividing patients by tumor HPV status (62.5% for HPV-negative tumors ν 20% for HPV-positive tumors), although HPV typing was available for only one third of our patient cohort.

We then constructed a risk stratification algorithm based on the previous data. Patients with HPV-positive tumors or with oropharyngeal primary tumors and no history of tobacco use were classified as low risk ($n = 61$). Those with HPV-negative tumors, nonoropharyngeal primaries, or significant tobacco use histories were classified as




Fig 3. Positive predictive value (PPV) of [18 F]fluorodeoxyglucose positron emission tomography (FDG-PET)/computed tomography (CT) by risk group. The PPV of FDG-PET/CT for primary and nodal disease combined is shown for patients with oropharyngeal (OP) and non-OP primaries, tobacco users and nonusers, and tumors negative or positive for human papillomavirus (HPV). The number of patients in each group is displayed below each data point.

high risk (n = 31). The rate of response to radiotherapy was greater in the low-risk group compared with the high-risk group (95.1% v 71.0%, respectively).

ROC curve analysis demonstrated that the accuracy of FDG-PET/CT was greater for the high-risk group compared with the low-risk group (Fig 4). This difference was most pronounced for the detection of residual nodal disease, where the area under the ROC curves was 50% greater for high-risk versus low-risk patients.

Finally, we compared the accuracy of both imaging modalities across the two risk groups (Table 2). The most pronounced benefit achieved by using FDG-PET/CT comes in detecting residual nodal disease for high-risk patients, where the positive predictive value of FDG-PET/CT is twice that of CT alone (75% v 37.5%, respectively). FDG-PET/CT also increases the specificity of response assessment




Fig 4. Risk-stratified [¹⁸F]fluorodeoxyglucose positron emission tomography (FDG-PET)/computed tomography (CT) receiver operating characteristic (ROC) curves. ROC curves for FDG-PET/CT are shown, separated by (A) primary and (B) nodal disease (high-risk patients = —; low-risk patients = ----). For comparison, the sensitivity/specificity of CT is plotted for high-risk (□) and low-risk (○) patients, as well.

Table 2. Accuracy of FDG-PET/CT by Risk Stratification

Parameter	Primary		Node	
	HR	LR	HR	LR
Sensitivity, %				
PET/CT	71.4	50.0	75.0	66.7
CT	83.3	75.0	100.0	80.0
NPV, %				
PET/CT	92.0	97.8	94.7	96.3
CT	95.7	98.0	100.0	96.7
Specificity, %				
PET/CT	100.0	82.7	84.2	57.9
CT	91.7	89.1	75.0	61.7
PPV, %				
PET/CT	100.0	16.7	75.0	14.3
CT	71.4	33.3	37.5	18.2

NOTE. Accuracy parameters are tabulated for FDG-PET/CT and CT alone, divided by primary and nodal sites, for both HR and LR patients.

Abbreviations: FDG, [¹⁸F]fluorodeoxyglucose; PET, positron emission tomography; CT, computed tomography; HR, high risk; LR, low risk; NPV, negative predictive value; PPV, positive predictive value.

compared with CT alone for high-risk patients, at the cost of slightly lower sensitivity and negative predictive value. For the low-risk group, FDG-PET/CT was inferior to CT alone for all metrics of response assessment accuracy.

DISCUSSION

Several retrospective series have investigated the utility of FDG-PET/CT imaging in HNSCC radiation response assessment.^{13-20,22} These series are confounded by limitations inherent to their retrospective design and by small sample size. A number of small prospective studies have been published using FDG-PET alone, without coregistered CT imaging, yielding inconsistent results.^{19,26,27} Two of these studies directly correlated imaging results with neck dissection findings; these demonstrated widely ranging staging accuracy (negative predictive values of 14% and 91.7%).^{19,27}

To our knowledge, the current study is the first to prospectively evaluate combined FDG-PET/CT for this indication. Our data confirm that FDG-PET/CT imaging provides little value over conventional CT imaging for assessment of radiation response in unselected patients with locally advanced head and neck cancer. Our analytic approach was to weight the cost of sensitivity and specificity equally to generate SUV_{max} thresholds with maximal overall accuracy. Doing so, FDG-PET/CT outperformed CT only in terms of specificity and positive predictive value. These characteristics are best suited for minimizing risk of unnecessary neck dissections. Most patients (72.7%) who underwent consolidative neck surgery on this protocol did so without clinical benefit, and there was no evidence that FDG-PET/CT findings influenced the rate or type of neck dissection performed (selective or complete). The increased specificity of FDG-PET/CT would seem suited to address this problem. Extrapolating from the data presented earlier, FDG-PET/CT could reduce the rate of false positives by approximately one third, eliminating false-positive imaging results for 12 patients in every 100 patients treated. Whether this benefit

outweighs a 67% relative increased risk of false-negative findings with FDG-PET/CT is worthy of debate. Our bias is to accept the risk of unnecessary consolidative dissections to avoid missing residual nodal disease. Given these data, then, we would not recommend routine use of FDG-PET/CT for radiation response assessment in unselected HNSCC patients.

Despite such findings, subset analysis revealed that FDG-PET/CT can provide benefit when applied in an individualized, risk-adjusted fashion. We initially surmised that selective use of FDG-PET/CT response assessment in patients at highest risk for recurrence would improve its diagnostic yield. Consistent with that premise, we demonstrated that FDG-PET/CT improved on standard CT response assessment for patients with HPV-negative tumors, nonoropharyngeal primaries, and histories of alcohol and tobacco use—patients who were at elevated risk for treatment failure. For these high-risk patients, nodal PET positivity after radiotherapy likely warrants surgical salvage (positive predictive value = 75%). A negative study should not necessarily preclude dissection of a neck where salvage is otherwise indicated because one in 20 of these patients would be in need of further therapy (negative predictive value = 95%).

The initial impetus to investigate HPV status in our patient population came from the observation that a history of heavy alcohol and tobacco use correlated with poor response to therapy on this trial. Tobacco use has been formally associated with tumor HPV negativity in a recent population-based study,²⁸ and tumor HPV negativity correlates strongly with poor response to radiotherapy²⁹ and shorter overall survival.^{29,30} We must caution that less than half of patients on this study had sufficient tissue available for analysis. This is a consequence of patients commonly presenting to our institution after a tissue diagnosis had been made at an outside facility. Nevertheless, our expanded subset analysis showed that other factors, namely nonoropharyngeal primary tumors and a history of significant tobacco use, also correlated with increased accuracy of FDG-PET/CT-based response assessment. We feel that the ability to triage patients based on primary tumor site and tobacco use is of great clinical relevance, given that HPV genotyping is not available universally.

Several limitations of this study merit attention. First, because this study was performed at a high-volume cancer center, the results may not be applicable to all treatment settings. The referral pattern at our institution is potentially different from other settings, where the proportion of high-risk patients and radiotherapy nonresponders may be greater. Our results suggest that FDG-PET/CT would be more accurate in such a scenario and may, therefore, deserve broader application in that setting. Second, this study was not designed to evaluate the impact of post-treatment interval on the accuracy of FDG-PET/CT, so this remains an open question.

The approach at our institution is to assess response 8 to 9 weeks after radiotherapy. This permits time for follow-up examinations by all providers, formal reading of imaging studies, and arrangement of necessary surgery within the time window preferred by our surgeons to minimize operative morbidity in the postradiation setting. Several groups have suggested that the accuracy of FDG-PET/CT scanning may increase with longer (10 to 12 weeks) postradiotherapy intervals.^{15,18,26,31} Other authors have described 8 weeks after radiotherapy as an appropriate cutoff interval to maximize accuracy.^{15,32} When our own data were stratified by post-treatment interval, we observed no

clear benefit to delaying imaging beyond 8 weeks (Appendix Figs A2 and A3, and Table A3, online only).

In conclusion, we prospectively demonstrate that FDG-PET/CT provides little value over CT alone in radiation response assessment for unselected patients with locally advanced HNSCC. Nonetheless, our data also suggest that FDG-PET/CT may improve assessment of treatment response in high-risk patients, such as those with HPV-negative disease. The currently enrolling Radiation Therapy Oncology Group 0522 phase III trial has incorporated a formal study of FDG-PET/CT imaging for treatment response assessment that closely mirrors the design of this current study. Confirmation of our results in the cooperative group setting would provide critical impetus to incorporate risk stratification strategies into FDG-PET/CT assessment of radiotherapy response in locally advanced head and neck cancer.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Although all authors completed the disclosure declaration, the following author(s) indicated a financial or other interest that is relevant to the subject matter under consideration in this article. Certain relationships marked with a "U" are those for which no compensation was received; those relationships marked with a "C" were compensated. For a detailed description of the disclosure categories, or for more information about ASCO's conflict of interest policy, please refer to the Author Disclosure Declaration and the Disclosures of Potential Conflicts of Interest section in Information for Contributors.

Employment or Leadership Position: None **Consultant or Advisory Role:** Homer A. Macapinlac, GE Healthcare (C), Radiology Corporation of America (C) **Stock Ownership:** None **Honoraria:** Homer A. Macapinlac, GE Healthcare, Siemens, Radiology Corporation of America **Research Funding:** Homer A. Macapinlac, GE Healthcare **Expert Testimony:** None **Other Remuneration:** None

AUTHOR CONTRIBUTIONS

Conception and design: Homer A. Macapinlac, K. Kian Ang, David L. Schwartz
Financial support: K. Kian Ang, Scott M. Lippman, David L. Schwartz
Administrative support: David L. Schwartz
Provision of study materials or patients: Lawrence E. Ginsberg, K. Kian Ang, K.S. Clifford Chao, Gregory M. Chronowski, William H. Morrison, David I. Rosenthal, Adam S. Garden, Scott M. Lippman, David L. Schwartz
Collection and assembly of data: Benjamin J. Moeller, Vishal Rana, Blake A. Cannon, Michelle D. Williams, Erich M. Sturgis, Lawrence E. Ginsberg, David L. Schwartz
Data analysis and interpretation: Benjamin J. Moeller, Michelle D. Williams, Erich M. Sturgis, Homer A. Macapinlac, J. Jack Lee, K. Kian Ang, Randal S. Weber, Scott M. Lippman, David L. Schwartz
Manuscript writing: Benjamin J. Moeller, Erich M. Sturgis, Homer A. Macapinlac, J. Jack Lee, David L. Schwartz
Final approval of manuscript: Benjamin J. Moeller, Vishal Rana, Blake A. Cannon, Michelle D. Williams, Erich M. Sturgis, Lawrence E. Ginsberg, Homer A. Macapinlac, J. Jack Lee, K. Kian Ang, K.S. Clifford Chao, Gregory M. Chronowski, Steven J. Frank, William H. Morrison, David I. Rosenthal, Randal S. Weber, Adam S. Garden, Scott M. Lippman, David L. Schwartz

REFERENCES

1. Denis F, Garaud P, Bardet E, et al: Final results of the 94-01 French Head and Neck Oncology and Radiotherapy Group randomized trial comparing radiotherapy alone with concomitant radiochemotherapy in advanced-stage oropharynx carcinoma. *J Clin Oncol* 22:69-76, 2004
2. Bonner JA, Harari PM, Giralt J, et al: Radiotherapy plus cetuximab for squamous-cell carcinoma of the head and neck. *N Engl J Med* 354:567-578, 2006
3. Jeremic B, Shibamoto Y, Milicic B, et al: Hyperfractionated radiation therapy with or without concurrent low-dose daily cisplatin in locally advanced squamous cell carcinoma of the head and neck: A prospective randomized trial. *J Clin Oncol* 18:1458-1464, 2000
4. Clayman GL, Johnson CJ 2nd, Morrison W, et al: The role of neck dissection after chemoradiotherapy for oropharyngeal cancer with advanced nodal disease. *Arch Otolaryngol Head Neck Surg* 127:135-139, 2001
5. Davidson BJ, Newkirk KA, Harter KW, et al: Complications from planned, posttreatment neck dissections. *Arch Otolaryngol Head Neck Surg* 125:401-405, 1999
6. Sassler AM, Esclamado RM, Wolf GT: Surgery after organ preservation therapy: Analysis of wound complications. *Arch Otolaryngol Head Neck Surg* 121:162-165, 1995
7. Lavertu P, Bonafede JP, Adelstein DJ, et al: Comparison of surgical complications after organ-preservation therapy in patients with stage III or IV squamous cell head and neck cancer. *Arch Otolaryngol Head Neck Surg* 124:401-406, 1998
8. Pellitteri PK, Ferlito A, Rinaldo A, et al: Planned neck dissection following chemoradiotherapy for advanced head and neck cancer: Is it necessary for all? *Head Neck* 28:166-175, 2006
9. McHam SA, Adelstein DJ, Rybicki LA, et al: Who merits a neck dissection after definitive chemoradiotherapy for N2-N3 squamous cell head and neck cancer? *Head Neck* 25:791-798, 2003
10. Peters LJ, Weber RS, Morrison WH, et al: Neck surgery in patients with primary oropharyngeal cancer treated by radiotherapy. *Head Neck* 18:552-559, 1996
11. Velázquez RA, McGuff HS, Sycamore D, et al: The role of computed tomographic scans in the management of the N-positive neck in head and neck squamous cell carcinoma after chemoradiotherapy. *Arch Otolaryngol Head Neck Surg* 130:74-77, 2004
12. Ojiri H, Mendenhall WM, Stringer SP, et al: Post-RT CT results as a predictive model for the necessity of planned post-RT neck dissection in patients with cervical metastatic disease from squamous cell carcinoma. *Int J Radiat Oncol Biol Phys* 52:420-428, 2002
13. Chen AY, Vilaseca I, Hudgins PA, et al: PET-CT vs contrast-enhanced CT: What is the role for each after chemoradiation for advanced oropharyngeal cancer? *Head Neck* 28:487-495, 2006
14. Connell CA, Corry J, Milner AD, et al: Clinical impact of, and prognostic stratification by, F-18 FDG PET/CT in head and neck mucosal squamous cell carcinoma. *Head Neck* 29:986-995, 2007
15. Gourin CG, Williams HT, Seabolt WN, et al: Utility of positron emission tomography-computed tomography in identification of residual nodal disease after chemoradiation for advanced head and neck cancer. *Laryngoscope* 116:705-710, 2006
16. Kubota K, Yokoyama J, Yamaguchi K, et al: FDG-PET delayed imaging for the detection of head and neck cancer recurrence after radio-chemotherapy: Comparison with MRI/CT. *Eur J Nucl Med Mol Imaging* 31:590-595, 2004
17. Nayak JV, Walvekar RR, Andrade RS, et al: Deferring planned neck dissection following chemoradiation for stage IV head and neck cancer: The utility of PET-CT. *Laryngoscope* 117:2129-2134, 2007
18. Porceddu SV, Jarmolowski E, Hicks RJ, et al: Utility of positron emission tomography for the detection of disease in residual neck nodes after (chemo)radiotherapy in head and neck cancer. *Head Neck* 27:175-181, 2005
19. Rogers JW, Greven KM, McGuirt WF, et al: Can post-RT neck dissection be omitted for patients with head-and-neck cancer who have a negative PET scan after definitive radiation therapy? *Int J Radiat Oncol Biol Phys* 58:694-697, 2004
20. Ware RE, Matthews JP, Hicks RJ, et al: Usefulness of fluorine-18 fluorodeoxyglucose positron emission tomography in patients with a residual structural abnormality after definitive treatment for squamous cell carcinoma of the head and neck. *Head Neck* 26:1008-1017, 2004
21. Wong RJ, Lin DT, Schoder H, et al: Diagnostic and prognostic value of [(18)F]fluorodeoxyglucose positron emission tomography for recurrent head and neck squamous cell carcinoma. *J Clin Oncol* 20:4199-4208, 2002
22. Yao M, Luo P, Hoffman HT, et al: Pathology and FDG PET correlation of residual lymph nodes in head and neck cancer after radiation treatment. *Am J Clin Oncol* 30:264-270, 2007
23. Hashibe M, Brennan P, Benhamou S, et al: Alcohol drinking in never users of tobacco, cigarette smoking in never drinkers, and the risk of head and neck cancer: Pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *J Natl Cancer Inst* 99:777-789, 2007
24. Dabaja B, Salehpour MR, Rosen I, et al: Intensity-modulated radiation therapy (IMRT) of cancers of the head and neck: Comparison of split-field and whole-field techniques. *Int J Radiat Oncol Biol Phys* 63:1000-1005, 2005
25. Garden AS, Morrison WH, Wong PF, et al: Disease-control rates following intensity-modulated radiation therapy for small primary oropharyngeal carcinoma. *Int J Radiat Oncol Biol Phys* 67:438-444, 2007
26. Greven KM, Williams DW 3rd, McGuirt WF Sr, et al: Serial positron emission tomography scans following radiation therapy of patients with head and neck cancer. *Head Neck* 23:942-946, 2001
27. Brkovich VS, Miller FR, Karnad AB, et al: The role of positron emission tomography scans in the management of the N-positive neck in head and neck squamous cell carcinoma after chemoradiotherapy. *Laryngoscope* 116:855-858, 2006
28. Gillison ML, D'Souza G, Westra W, et al: Distinct risk factor profiles for human papillomavirus type 16-positive and human papillomavirus type 16-negative head and neck cancers. *J Natl Cancer Inst* 100:407-420, 2008
29. Fakhry C, Westra WH, Li S, et al: Improved survival of patients with human papillomavirus-positive head and neck squamous cell carcinoma in a prospective clinical trial. *J Natl Cancer Inst* 100:261-269, 2008
30. Ragin CC, Taioli E: Survival of squamous cell carcinoma of the head and neck in relation to human papillomavirus infection: Review and meta-analysis. *Int J Cancer* 121:1813-1820, 2007
31. Yao M, Smith RB, Graham MM, et al: The role of FDG PET in management of neck metastasis from head-and-neck cancer after definitive radiation treatment. *Int J Radiat Oncol Biol Phys* 63:991-999, 2005
32. Andrade RS, Heron DE, Degirmenci B, et al: Posttreatment assessment of response using FDG-PET/CT for patients treated with definitive radiation therapy for head and neck cancers. *Int J Radiat Oncol Biol Phys* 65:1315-1322, 2006